

Rat Monoclonal Anti-human Seprase Antibody D28

PRODUCT INFORMATION

Catalog Number:	MABS1002
Hybridoma Clone:	D28
Lot Number:	A-002
Quantity:	0.1 mg
Concentration:	1.0 mg/mL
Antibody Type:	Rat IgG2a
Formulation:	0.1 mg antibody in protein-free hybridoma medium, PBS, pH 7.4, and 50% glycerol.
Storage:	- 20° C
Specificity:	Human seprase (also called fibroblast activation protein alpha, FAP α)
Immunogen:	Seprase isolated from human termed placenta
Applications:	Immunohistochemistry, Immunocytochemistry, Immunofluorescence, Flow cytometry, ELISA, Immunocapture, Immunoprecipitation, Flow Cytometry and Western blotting.
Method of protein determination:	SDS-PAGE analysis showing greater than 99% protein being IgG heavy chain at 55-kDa and light chain at 25-kDa and Bradford method.
Method of activity determination:	Immuno-capture of recombinant antigen produced by 293-EBNA human kidney cells.

DESCRIPTION

This antibody is produced from the D28 hybridoma cell line derived from fusion of rat myeloma Y3 cells and spleen cells of an immunized Sprague-Dawley rat (Pineiro-Sanchez et al., 1997). Seprase, also called fibroblast activation protein alpha (FAP α) [Gene ID: 2191; Accession#: NP_004451] is a 170-kDa homodimeric integral membrane gelatinase belonging to the type II transmembrane serine protease family. It has a non-classical serine catalytic site and exhibits dipeptidyl dipeptidase and gelatinase activities. Seprase is selectively expressed in various cell types from cancerous tissues as well as tissues in the early stages of wound healing (Chen et al., 2003). This protein is thought to be involved in tumor invasion, angiogenesis and metastasis as well as embryo development and tissue repair. Recently, truncated and active forms of seprase have been detected in human malignant tumors (Chen et al., 2006) and plasma (Lee et al., 2006).

PREPARATION

MAb D28 was produced by D28 hybridoma cells in protein-free medium using CELLLine CL 1000 Disposable Bioreactors (INTEGRA Biosciences) that includes a 10 kDa semi-permeable cellulose acetate membrane to exclude small molecules. Supernatant from the cell compartment was cleared by spinning at 10,000rpm. Approximate 5 mg/mL of antibody was obtained and diluted to 2.0 mg/mL using PBS, pH 7.4. Equal volume of glycerol was added to the antibody solution to make the final concentration of 1.0 mg/mL.

METHOD OF ACTIVITY DETERMINATION

The activity of the antibody was determined using an immunocapture assay. Briefly, anti-rat antibody was coated on a plate and 50 μ g/mL of target rat monoclonal antibody was added. Native seprase from human tumor cells or recombinant human seprase was then added. The prolyl dipeptidase and serine gelatinase activities of the captured seprase were measured using seprase substrates Gly-Pro-p-Nitroanilide (sigma) and DQ gelatin (Invitrogen), respectively. Alternatively, antigen captured by the target antibody was detected using biotinylated anti-seprase antibody D8, followed by a colorimetric assay.

METHOD OF PROTEIN DETERMINATION

Protein concentration was determined using SDS-PAGE under reducing and denaturing conditions. Antibody protein was determined for its identity by SDS-PAGE analysis that shows greater than 99% of total protein being IgG heavy chain at 55-kDa and light chain at 25-kDa (Figure 1). The total protein in the preparation was measured with Bradford protein assay using Quick Start Bradford Dye Reagent (Bio-Rad); serially diluted BSA samples were used as standards.




Figure 1. SDS-PAGE analysis of mAb D28 under denaturing and reducing conditions. 5 μ g/lane.

STORAGE

The antibody may be stored at -20° C for one year in its original formulation. Additionally, antibody diluted with 1% BSA in PBS may be stored at 2° to 8° C for up to 1 month without detectable loss of activity. **Avoid repeated freeze-thaw cycles of the diluted antibody.**

SPECIFICITY

This antibody recognizes human full-length seprase and recombinant seprase excluding the cytoplasmic (amino acids 1-6) and transmembrane domains of seprase (amino acids 7-26). This antibody recognizes an epitope that is different from the binding sites of mAb D8, mAb D43 and mAb E97. In Western blotting, it does not recognize mouse seprase (Chen et al., 2006) and human DPPIV, a homolog of seprase (Gherzi et al., 2002); in immunoprecipitation, it does not detect human DPPIV (Gherzi et al., 2006). Furthermore, this antibody does not block seprase peptidase activity against Gly-Pro-pNA (Gherzi et al., 2002; Gherzi et al., 2006); it does inhibit the gelatinase activity of seprase.

APPLICATIONS

Immunohistochemistry – Monoclonal anti-human seprase antibody D28 may be used to detect human seprase via immunohistochemistry techniques. Using an antibody concentration of 1 to 5 µg/mL, human seprase has been identified in paraffin-embedded tissue sections (Gherzi et al., 2006; Iwasa et al., 2005). Antigen retrieval may be achieved by heating in an autoclave in 0.01M citrate buffer (pH6.0). This antibody has also been used to detect human seprase in frozen tissue sections (Gherzi et al., 2002).

Immunocytochemistry and Immunofluorescence – This antibody is also effective for direct and indirect immunofluorescence staining of human seprase at cell surfaces. Human seprase concentrated at cell protrusions of tumor and endothelial cells as well as fibroblasts was identified via fluorescence and confocal microscopy using an antibody concentration of 1 to 10 µg/mL (Gherzi et al., 2002; Gherzi et al., 2006).

ELISA – This antibody can be used in ELISA at a concentration of 5 to 50 µg/mL to detect purified seprase coated on plastic plates (Nakahara et al., 1996).

Immuno-capture – This antibody has been employed to capture native seprase and recombinant seprase in their active forms (Chen et al., 2006; Gherzi et al., 2002). In one assay performed (Figure 2), the antibody is linked to a solid support to capture seprase in cell lysates. Then the prolyl dipeptidase and serine gelatinase activities of seprase are measured using seprase substrates Gly-Pro-p-Nitroanilide (sigma) and DQ gelatin (Invitrogen), respectively.




Figure 2. Immunocapture analysis of seprase from human melanoma LOX cell lysate. Cell lysate was applied to plastic wells coated with mAb D28 and the captured seprase was detected by measuring its prolyl dipeptidase and serine gelatinase activities as described (Gherzi et al., 2002). PBS was used as a negative control for the cell lysate. The experiment was performed in triplicate and the values are the mean ±SD.

Immunoprecipitation – This antibody has been conjugated to agarose beads to successfully immuno-precipitate human seprase from human tumor tissue lysates (Chen et al., 2006) and lysates from different cells in culture (Gherzi et al., 2002; Gherzi et al., 2006; Mueller et al., 1999; Monsky et al., 1994). This antibody may also work indirectly for immunoprecipitation by using anti-rat IgG antibody conjugated beads.

Flow Cytometry- This antibody has been used to sort malignant melanoma LOX cells with high seprase expression on the cell surface (Monsky et al., 1994).

Western Blotting-This antibody can be used at 1 to 10 µg/mL with the appropriate secondary reagents to detect human seprase in Western blotting (Pineiro-Sanchez et al., 1997; Monsky et al., 1994). This antibody recognizes both seprase 170-kDa dimers and the dissociated 97-kDa monomers.




Figure 3. Western blotting analysis of seprase using mAb D28. Purified seprase from melanoma LOX cell was subjected to SDS-PAGE under non-denaturing (N) or denaturing (D) conditions, followed by immunoblotting using mAb D28 as described (Pineiro-Sanchez et al., 1997). MAb D28 recognizes both seprase 170-kDa dimer and the 97-kDa subunit.

REFERENCES

- Chen,D., Kennedy,A., Wang,J.Y., Zeng,W., Zhao,Q., Pearl,M., Zhang,M., Suo,Z., Nesland,J.M., Qiao,Y., Ng,A.K., Hirashima,N., Yamane,T., Mori,Y., Mitsumata,M., Ghersi,G., and Chen,W.-T. (2006). Activation of EDTA-Resistant Gelatinases in Malignant Human Tumors. *Cancer Res.* 66, 9977-9985.
- Chen,W.-T., Kelly,T., and Ghersi,G. (2003). DPPIV, seprase and related serine peptidases in multiple cellular functions. In *Cell surface proteases*, S.Zucker and W.-T.Chen, eds. (San Diego, CA.: Academic Press), pp. 207-232.
- Ghersi,G., Dong,H., Goldstein,L.A., Yeh,Y., Hakkinen,L., Larjava,H.S., and Chen,W.-T. (2002). Regulation of fibroblast migration on collagenous matrix by a cell surface peptidase complex. *J. Biol. Chem.* 277, 29231-29241.
- Ghersi,G., Zhao,Q., Salamone,M., Yeh,Y., Zucker,S., and Chen,W.-T. (2006). The protease complex consisting of dipeptidyl peptidase IV and seprase plays a role in the migration and invasion of human endothelial cells in collagenous matrices. *Cancer Res* 66, 4652-4661.
- Iwasa,S., Okada,K., Chen,W.-T., Jin,X., Yamane,T., Ooi,A., and Mitsumata,M. (2005). 'Increased expression of seprase, a membrane-type serine protease, is associated with lymph node metastasis in human colorectal cancer'. *Cancer Lett.* 227, 229-236.
- Lee,K.N., Jackson,K.W., Christiansen,V.J., Lee,C.S., Chun,J.G., and McKee,P.A. (2006). Antiplasmin-cleaving enzyme is a soluble form of fibroblast activation protein. *Blood* 107, 1397-1404.
- Monsky,W.L., Lin,C.-Y., Aoyama,A., Kelly,T., Mueller,S.C., Akiyama,S.K., and Chen,W.-T. (1994). A potential marker protease of invasiveness, seprase, is localized on invadopodia of human malignant melanoma cells. *Cancer Res.* 54, 5702-5710.
- Mueller,S.C., Ghersi,G., Akiyama,S.K., Sang,Q.X., Howard,L., Pineiro-Sanchez,M., Nakahara,H., Yeh,Y., and Chen,W.-T. (1999). A novel protease-docking function of integrin at invadopodia. *J. Biol. Chem.* 274, 24947-24952.
- Nakahara,H., Nomizu,M., Akiyama,S.K., Yamada,Y., Yeh,Y., and Chen,W.-T. (1996). A mechanism for regulation of melanoma invasion. Ligation of alpha6beta1 integrin by laminin G peptides. *J. Biol. Chem.* 271, 27221-27224.
- Pineiro-Sanchez,M.L., Goldstein,L.A., Dodt,J., Howard,L., Yeh,Y., Tran,H., Argraves,W.S., and Chen,W.-T. (1997). Identification of the 170-kDa melanoma membrane-bound gelatinase (seprase) as a serine integral membrane protease. *J. Biol. Chem.* 272, 7595-7601; Correction (1998) *J. Biol. Chem.* 273, 13366.

For Research Use Only. Not For Diagnostic or Therapeutic Use.

Conditions: Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Vitalex Inc is strictly prohibited.